

Chapter 3 – ADVERSE EFFECTS OF TOBACCO USE IN DEPLOYED MILITARY UNITS

by

Merle PARMAK, PhD (EST) and Marten MEIJER, PhD (NLD)

ABSTRACT

Although research reveals that smoking prevalence has in general stabilized or is even decreasing among military personnel, this trend does not ultimately apply. Being young, being deployed, or being a member of Army personnel, for instance, is proven to increase the risk of being or beginning to be a tobacco user. Usually there are not immediate links emphasised between tobacco habits and the serious health-related consequences during the service period because of the long time lag between tobacco use and its consequences. With some exceptions, the impact of smoking on military performance is defined rather indirectly than directly. However, findings in the Estonian military sample (n = 135) indicate that an increase in smoking behaviour while on deployment not only corresponds with poorer psychological well-being and general health, but also with being forced to stay away from duty because of physical aches. These results can be taken as indicators that smoking behaviour decreases fitness for military operations and should be targeted by performance enhancement activities. The role of the military culture of smoking behaviour and the arguments for an effective strategy for tobacco use cessation among military personnel are discussed.

3.1 TRENDS IN TOBACCO USE

Considering the direct and indirect costs related to consequences of tobacco habits, high importance has been attributed to this behaviour in health policies. Tobacco use is known as the single largest cause of preventable death in the world today, killing a third to a half of all users. Projecting into the future, the total tobacco-attributable deaths will account for almost 10% of all deaths worldwide in 2030 [1] In the WHO European Region, smoking is blamed for about 18.6 million years of life lost [2].

The financial costs caused by tobacco-related illnesses and medical care are remarkable. The direct and indirect costs of smoking in the EU, for instance, were estimated to range from 1.04% to 1.39% of the EU Gross Domestic Product, exceeding even 3% of it in some new member countries [3].

Some pessimistic prognoses show that the worldwide number of smokers continues to increase [4] and that the deaths caused by tobacco will double over the next few decades [5]. However, the trends of tobacco-related habits in the Western world are constantly decreasing. According to the latest health surveys in the US and in Europe, smoking prevalence among men and women has in general stabilized or is even decreasing. For example, in the US, the past month use of tobacco products was 29.4 % in 2005, while it was 30.4% in 2002 [6]; in the WHO European Region, smoking prevalence was estimated around 28.6% in 2005 but 28.8% in 2002. Falling death rates due to tobacco-related illnesses imply that trends in smoking prevalence have been curbed at least since the early eighties [7].

Findings from Western military surveys show similar trends and indicate overall declines in smoking as well. For example, in the total military population, the prevalence of any smoking in the US declined from 51.0% in 1980 to 32.2% in 2005 [8]. In the Canadian Forces, everyday smoking has decreased from 24% in 2000 to 20% in 2004; the latter figure is declared to be even lower than in the civilian population [9]. However, low smoking rates in the armed forces are not the case in every country, and cannot be generalized to all tobacco products or age groups [9]. In Estonia, for instance, we can observe the same smoking prevalence as elsewhere in Europe (i.e., 27.8%) in the general population [10], but the prevalence

is as high as 41.1% in the military population [11]. Recent findings also reveal that military personnel are more than twice as likely as civilians to use smokeless tobacco [12] and also indicate an increasing rate of tobacco use among young military members [13].

Inside the military, tobacco use rates and initiation or cessation related aspects are widely explored, especially in relation to the extra stress or excessive boredom military personnel might have experienced on duty. Being deployed has been found to be associated with higher rates of cigarette use [14]. There are an increasing number of regular smokers (including relapse and new initiation) of approximately 10% as well as an increase in daily consumption from an average of 15 cigarettes to 21 cigarettes [15]. The main reasons for increased smoking during deployments that have been cited are:

- 1) Stress, boredom, anxiety, and sleep deprivation;
- 2) Lack of alternate activities and privileges;
- 3) The perception that dangers in the field trump the negative health impact of smoking; and
- 4) A permissive military culture toward tobacco use [16].

With respect to managing stress, however, the research findings indicate that tobacco use is more likely to perpetuate a stress response rather than to suppress it, and that nicotine consumers are overall less effective in dealing with combat stress [17]. Unfortunately, not much can be found about the lastingness of post-deployment changes in smoking behaviour. One survey where the persistence of this behaviour is described indicates noticeable differences among sub-groups: a larger percentage of Army personnel began or increased their cigarette smoking one year after having deployed than stopped or reduced, whereas the opposite was reported for the other services [9].

3.2 TOBACCO USE AND MILITARY FITNESS

Clinical studies have reported that cigarette use is associated with a lower functional status [18] and a lower exercise tolerance among young adult people [19];[20]. Smoking has also been found to be a consistent and strong predictor of the lack of fitness for military duty, operationalized, e.g., in measures of physical health, mental health, substance abuse, and legal problems and of the occurrence of medical problems in training. It is even suggested that smoking be considered as a negative marker of readiness and be included in the services' fitness evaluations [21];[22]. Considering the frequency of injury incidents in training [23] and in infantry duties [24] related to cigarette smoking, it has been cited as an independent risk factor for both men and women [25] Similar findings about the harmful effect of cigarette smoking on physical fitness and readiness are described among U.S. Navy personnel [26].

It is shown that cigarette smoking adversely impacts troop readiness with increasing time off from duty [27], leading to poorer visual acuity [28], and together the exposure to fine dust being possibly related with the acute eosinophilic pneumonia [29]. However, the decreased fitness during a military exercise, even in harsh environments and in combination with poor dietary habits [30] is not clearly and explicitly identified. Study results remain controversial about the harmful effects of smokeless tobacco as well. On the one hand, there are results showing that using smokeless tobacco is an independent risk factor for injury proneness [31], that it has a detrimental effect on visuo-motor performance [32], and that it is associated with hypercholesterolemia [33] and higher blood pressure [34]. On the other hand, results indicate that even long-term use of smokeless tobacco does not significantly influence exercise capacity [35]. However, even while physical performance may remain unaffected, there is an increased risk of all kinds of oral problems for users [36];[37] and a negative effect on performance caused by deprivation symptoms, such as increased reaction time, self-rated withdrawal and decreased heart rate [38].

Military fitness is not only about physical health and hardiness. Several studies have reported the association between cigarette smoking and psychiatric illnesses. For currently enlisted personnel, smoking

is found to be one of the factors predictive of hospitalization for mental health disorders [39]. Regarding psychiatric illness research, in the target group with Post-Traumatic Stress Disorder (PTSD) or major depression, there is a higher prevalence of smoking. It refers to possible self-medication caused by the alleviating effect of nicotine on some symptoms like arousal, numbness, or detachment, which are related to these disorders [40]. The fact that poor mental health relates to failures in smoking cessation [41] indicates that for those people, quitting is even more difficult than for healthy people. It has also been shown that the overall quality of life among veterans is affected by poor health behaviours, even after controlling for the impact of co-morbid medical conditions [42]. Taking a closer look at the average level of self care among PTSD veterans, one can also observe quite low frequencies of preventive health behaviours and increased risks for non-fatal strokes and myocardial infarctions [43].

3.3 INTERVENTION OPPORTUNITIES

Often there is more than one health risk behaviour or kind of substance in use involved simultaneously [44];[45] and it is difficult to detect which of them is responsible for the given disease or harm. The fact that usually several risky behaviours are concurrently present is observable among teenagers in the civil population [46];[47] as well as among the adult population in a military environment (e.g., high-risk drinkers use seatbelts less frequently, are more likely to exceed speed limits while driving and smoke more than 20 cigarettes per day). Therefore, intervention programs should be implemented for all those behaviours (safe driving habits, smoking cessation, high-risk drinking) and to be tailored to the specific needs of the group at highest risk [48]. However, when expecting a positive change, one should be aware that people do not alter several behaviours at the same time and efforts to modify one kind of unhealthy behaviour into a healthy one will not necessarily affect other risky behaviours [49].

The struggle for a healthy lifestyle in the military is far from hopeless as tobacco interventions aimed at smoking cessation have proven to be effective among veterans [50] as well as active duty military personnel [51]. Others have provided an exhaustive list of evidence-based practices of tobacco-control programs and activities are described in depth [52], and clinical treatment approaches [53]. Still, more needs to be done to change the military culture, which has been invoked as a kind of excuse for tobacco consumption (i.e., a means for enhancing comfort or as a morale booster) in almost every article or health report dealing with this population. It has been proven that social influence encourages tobacco use [54], and role models of smoking behaviour in the military are strongly associated with the initiation and resumption of smoking, even after adjusting for other known risk factors [55]. Consequently, intervening with empty slogans or vague efforts is ineffective. Without trying to modify the organisational culture, a persistent change in behaviour can hardly be reached.

As an additional hidden menace to the culture of the Armed Forces, military personnel form an attractive market segment for tobacco producers. Manufacturers' business interests are expressed in manipulative messages, openly directed to military members with high effectiveness [56]. This should be taken into account when trying to protect military members from (re)starting tobacco use and when elaborating the strategy for tobacco use cessation. To reduce existing perverse incentives that lead to increased tobacco consumption, an effective tobacco control policy in the Armed Forces requires explicit implementation instructions and high-level organisational support [57]. Extra attention should be paid to formulate segment-specific messages for military members that counteract effectively with industry messages.

3.4 RECENT FINDINGS

A survey [58] was conducted among two rotations of Estonian soldiers deployed into Afghanistan to figure out the change in their perceived general health and smoking behaviour during the first three months of deployment as well as to explore the relations between declared changes in behaviour and the level of psychological well-being reported. The use of other tobacco products like snuff was not explored in this research. All respondents were white males.

Soldiers (n = 135) were asked if they had noticed a change in their:

- 1) General health;
- 2) Seeking help from a physician;
- 3) Being forced to stay away from duty because of aches;
- 4) Smoking behaviour; and
- 5) Frequency of physical fitness training compared with the period before deployment.

The questions of being a smoker versus non-smoker, and – if yes – the number of cigarettes smoked per day were not explicitly asked for. Nevertheless, the smokers could be distinguished from the non-smokers through item x shown below. Participants had three choices to answer an item indicating a change:

- 1) Negative change;
- 2) Positive change; or
- 3) No changes in the health related aspect considered.

Examples of items:

x. Compared with the period before deployment my smoking behaviour did ...

- 1) *Increase;*
- 2) *Decrease; and*
- 3) *Not change (did not start if non-smoker).*

xx. Compared with the period before deployment my general health is ...

- 1) *Worse;*
- 2) *Better; and*
- 3) *Unchanged.*

To assess psychological well-being, we asked them to fill out the well-being questionnaire World Health Organization-Five Well-Being Index (WHO-5) [59]. For both rotations, a survey was administered in the middle of their deployment (being May and August 2008, respectively) on their way to Rest and Recuperation¹ in Estonia.

Most participants declared no changes in assessed health related aspects in the middle of deployment (Figure 3-1). Some negative trend was found in all health related aspects but the second highest for Smoking behaviour (n = 29) after frequency of Fitness training (n = 40). In terms of positive change, we observed the highest change for the General health which was evaluated more positively (better general health) at the mid deployment as compared with the period before deployment.

¹ Rest and Recuperation stands for a short break in the middle of deployment, in which troops visit home or stay in a safe area near the theatre of operations.




Figure 3-1: Dynamics of Self-Reported Changes in Health Related Aspects in the Middle of Deployment.

The odds ratios in change (positive against negative changes) are also very informative. If the number is higher than 1 then there are proportionally more positive changes than negative ones; and if the number is lower than one, then there are more negative changes as compared to the positive ones. The odds ratio for General health is 3.5; this means that there are approximately 3.5 times more positive changes than negative ones. On the contrary, the odds ratio for Smoking behaviour is 0.30; in other words, there are about three times more negative changes than positive ones.

The matrix presented in Table 3-1 shows correlations between health-related aspects measured among all respondents (n = 135). Higher scores on the scales indicate a positive change: better psychological well-being, better general health, fewer visits to a physician, less excessive aches, decreased smoking, and more physical fitness training. Hence, higher positive correlations have a positive connotation. Results reveal that the correlations between Smoking behaviour and Psychological well-being (r = .31), General health (r = .36), and Aches (r = .28) are positive and significant (p < .01). Notwithstanding the fact that correlations do not allow for any causal relationship between the variables, the results show that the observed negative changes in tobacco use (thus more smoking – see Figure 3-1) and in general health is correlated with poorer psychological well-being during the first three months of deployment.

Table 3-1: Correlations Between Psychological Well-Being and Health Aspects.

	Psy.WB	Gen.Hlth	Seek.Phy	Ex. Aches	Smoking
Psychological WB	–				
General Health	.33*	–			
Seeking a Physician	.11	.38*	–		
Excessive Aches	.25	.31*	.14	–	
Smoking Behaviour	.31*	.36*	.05	.28*	–
Fitness Training	.04	.37*	.21	.14	.21

Note. Psy.WB – Psychological Well-Being, Gen.Hlth – General Health, Seek.Phy – Seeking a Physician, Ex.Aches – Excessive Aches, Smoking – Smoking Behaviour

* $p < .01$

3.5 DISCUSSION

Findings from empirical part of the review concur with the line of previous research indicating that on deployments smoking behaviours is increase. Together with the perceived state of general health, smoking impacts soldiers’ psychological well-being. It is shown that soldiers who experience problems with their general health and whose smoking behaviour become more frequent are more vulnerable to mental distress. On the other hand smoking behaviour itself is predicted by the state of general health and presence of aches. It seems to be a closed circle of afore mentioned self medication where one problem is cured with the other and no easy solution is available. Military personnel on deployments are relatively young. Considering the remarkable time lag between tobacco use and its consequences, we might expect the harm to become more disturbing among older soldiers and among retired military members and veterans in terms of restricting their everyday activities, impairing quality of life, and reducing life expectancy.

In contrast with alcohol consumption or risky driving, there is not such an immediate impact of tobacco habits and the harm caused by those bad habits, such as increased death, premature deceases, serious injury rates or severe diseases found during the active duty service period. In active duty the impact of tobacco use on general health and specifically on military performance can be defined rather more indirectly. However, this indirect impact (e.g., injury proneness, deprivation symptoms, higher blood pressure, impaired vision acuity) of tobacco related behaviour may still decrease troops’ fitness for military operations and should be the target of performance enhancement activities. Based on research recommended interventions include:

- 1) Working out an intervention programs tailored to the specific needs of the group at highest risk;
- 2) Elaborating the strategy to modify the military culture which encourage tobacco use;
- 3) Developing an effective tobacco control policy in the Armed Forces with explicit implementation instructions and high-level organisational support; and
- 4) Formulating segment-specific messages for military members that are able to counteract effectively with messages from tobacco industries.

Research has shown that, instead of pointing to the manipulations by the tobacco industry or to the unhealthy effects of tobacco use, there are four promising themes for tobacco control efforts in the military. Messages to this population should emphasise that:

- 1) Smoking decreases one’s ability to positively influence others;

- 2) Smoking increases the chance that a military member will be discharged from the military prematurely;
- 3) Smoking lowers the readiness to fight and to win wars; and
- 4) Smokers are not as productive as other military personnel [60].

3.6 REFERENCES

- [1] World Health Statistics 2008. Part 1 Ten highlights in health statistics. Reducing deaths from tobacco.
- [2] WHO The world health report (2002). Reducing Risks, Promoting Healthy Life (<http://www.who.int/whr>).
- [3] WHO Regional Office for Europe. The European Tobacco Control Report 2007 (<http://www.euro.who.int>).
- [4] Guindon, G. E. and Boisclair, D. (2003) Past, current and future trends in tobacco use. Washington, DC, World Bank (<http://www1.worldbank.org>).
- [5] WHO Report on the Global Tobacco Epidemic, 2008: The MPOWER package, Geneva, World Health Organization, 2008 (<http://www.who.int>).
- [6] Substance Abuse and Mental Health Services Administration. (2006). Results from the 2005 National Survey on Drug Use and Health: National Findings (Office of Applied Studies, NSDUH Series H-30, DHHS Publication No. SMA 06-4194). Rockville, MD. (<http://www.samhsa.gov>).
- [7] WHO Regional Office for Europe (2007). The European tobacco control report 2007 (<http://www.euro.who.int>).
- [8] RTI International (2006) 2005 Department of Defence Survey of health Related Behaviours among Active Duty Military Personnel. RTI/7841/106-FR.
- [9] Directorate of Force Health Protection CF Health Services Group (2005). Canadian Forces Health and Lifestyle Information Survey 2004 Regular Force Report A-MD-015-FHP/AF-002.
- [10] Estonian National Institute for Health Development. Health behaviour among the Estonian adult population 2006 (<http://www2.tai.ee/uuringud>).
- [11] Estonian National Institute for Health Development (2008). Prevalence of smoking behaviour in Estonian Defence Forces. (preliminary data).
- [12] Peterson, A.L., Severson, H.H., Andrews, J.A., Gott, S.P., Cigrang, J.A., Gordon, J.S., Hunter, C.M. and Martin, G.C. (2007). Smokeless tobacco use in military personnel. *Military Medicine*, 172, 1300-1305.
- [13] Nelson, J.P. and Pederson, L.L. (2008). *Nicotine & Tobacco Research*, 10, 775-790.
- [14] Federman, B.E., Bray, M.B. and Kroutil, L.A. (2000). Relationships between substance use and recent deployments among women and men in the military. *Military Psychology*, 12, 205-220.
- [15] Boos, J.C. and Coft, A.F. (2004). Smoking rates in the staff of a military field hospital before and after wartime deployment. *Journal of the Royal Society of Medicine*, 97, 20-22.

- [16] Poston, W.S.C., Taylor, J.E., Hoffman, K.M., Peterson, A.L., Lando, H.A., Shelton, S. and Haddock, C.K. (2008). Smoking and Deployment: Perspectives of Junior-Enlisted U.S. Air Force and U.S. Army Personnel and Their Supervisors. *Military Medicine*, 173, 441-447.
- [17] Stein, R.J., Pyle, S.A., Haddock, C.K., Poston, W.S.C., Bray, R. and Williams, J. (2008). Reported stress and its relationship to tobacco use among U.S. military personnel. *Military Medicine*, 173, 271-277.
- [18] Eriksen, W., Natwig, B., Rutle, O. and Bruusgard, D. (1999). Smoking and functional status of young adults. *Scandinavian Journal of Primary Health Care*, 17, 174-179.
- [19] Papathanasiou, G., Georgakopoulos, D., Georgoudis, G., Spyropoulos, P., Perrea, D. and Evangelou, A. (2007). Effects of chronic smoking on exercise tolerance and on heart rate-systolic blood pressure product in young healthy adults. *European Journal of Cardiovascular Prevention and Rehabilitation*, 14, 646-652.
- [20] Kimura, Y., Nakamoto, Y., Shitama, H., Ohmine, S., Ide, M. and Hashisuka, K. (2007). Influence of moderate smoking on physical fitness and local muscle oxygenation profile during incremental exercise. *Journal of UOEH*, 29, 149-158.
- [21] Haddock, C.K., Pyle, S.A., Poston, W.S.C., Bray, R.M. and Stein, R.J. (2007). Smoking and Body Weight as Markers of Fitness for Duty among U.S. Military Personnel. *Military Medicine*, 172, 527-532.
- [22] Snoddy, R.O. and Henderson, J.M. (1994). Predictors on basic infantry training success. *Military Medicine*, 159, 616-622.
- [23] Reynolds, K.L., Heckel, H.A., Witt, C.E., Martin, J.W., Polland, J.A., Knapik, J.J. and Jones, B.H. (1994). Cigarette smoking, physical fitness, and injuries in infantry soldiers. *American Journal of Preventive Medicine*, 10, 145-150.
- [24] Reynolds, K.L., White, J.S., Knapik, J.J., Witt, C.E. and Amoroso, P.J. (1999). Injuries and risk factors in a 100-mile (161-km) infantry road march. *Preventive Medicine*, 28, 167-173.
- [25] Knapik, J.J., Sharp, M.A., Canham-Chervak, M., Hauret, K., Patton, J.F. and Jones, B.H. (2001). Risk factors for training-related injuries among men and women in basic combat training. *Medicine and Science in Sports Exercise*, 33, 946-954.
- [26] Van Orden, K.F. and Nice, D.S. (2008). National security interests at the naval health research center. In A.D. Mangelsdorff (Ed.), *Psychology in the service of national security* (pp. 55-70). Washington, DC: American Psychological Association.
- [27] Zadoo, V., Fengler, S. and Catteson, M. (1993). The effects of alcohol and tobacco use on troop readiness. *Military Medicine*, 158, 480-484.
- [28] Sharma, R.R. (2009). Visual effect of long-term active smoking: Are air-crew flying NVG aided missions at a disadvantage? Paper presented at 57-th International Congress of Aviation and Space Medicine, Zagreb, 2009.
- [29] Shorr, A.F., Scoville, S.L., Cersovsky, S.B., Shanks, G.D., Ockenhouse, C.F., Smoak, B.L., Carr, W.W. and Petrucci, B.P. (2004). Acute eosinophilic pneumonia among US Military personnel deployed in or near Iraq. *The Journal Of The American Medical Association*, 292, 2997-3005.

- [30] Defence Science and Technology Organisation Victoria (Australia) Aeronautical and Maritime Research Lab (2002). The Effect of Consumption of Australian Combat Rations on Military Personnel after a Medium-Term Field Exercise (<http://handle.dtic.mil/100.2/ADA399668>).
- [31] Heir, T. and Eide, G. (1997). Injury proneness undergoing a physical training programme: smokeless tobacco use, higher age, and low levels of physical fitness are risk factors. *Scandinavian Journal of Medicine & Science in Sports*, 7, 304-311.
- [32] Contreras-Vidal, J.L., Van den Heuvel, C.E., Teulings, H.L. and Stelmach, G.E. (1999). Visuo-motor adaptation in smokeless tobacco users. *Nicotine & Tobacco Research*, 1, 219-227.
- [33] Tucker, L.A. (1989). Use of smokeless tobacco, cigarette smoking, and hypercholesterolemia. *American Journal of Public Health*, 79, 1048-1050.
- [34] Bolinder, G. and De Faire, U. (1998). Ambulatory 24-h blood pressure monitoring in healthy, middle-aged smokeless tobacco users, smokers and non-tobacco users. *American Journal of Hypertension*, 11, 1153-1163.
- [35] Bolinder, G., Noren, A., Wahren, J. and De Faire, U. (1997). Long-term use of smokeless tobacco and physical performance in middle-aged men. *European Journal of Clinical Investigation*, 27, 427-433.
- [36] Robertson, P.B., De Rouen, T.A., Ernster, V., Grady, D., Greene, J., Mancl, L., McDonald, D. and Wash, M.M. (1995). Smokeless tobacco use: how it affects the performance of major league baseball players. *Journal of the American Dental Association*, 126, 1115-1121.
- [37] Robertson, P.B., Walsh, M.M. and Greene, J.C. (1997). Oral effects of smokeless tobacco use by professional baseball players. *Advances in Dental Research*, 11, 307-312.
- [38] Keenan, R.M., Hatsukami, D.K. and Anton, D.J. (1989). The effects of short-term smokeless tobacco deprivation on performance. *Psychopharmacology*, 98, 126-130.
- [39] Booth-Kevley, S. and Larson, G.E. (2005). Predictors of Psychiatric Hospitalization in the Navy. *Military Medicine*, 170, 87-93.
- [40] RAND Centre for Military Health Policy Research (2008). Invisible Wounds of War (<http://www.rand.org>).
- [41] Hemmingsson, T., Kriebel, D., Tynelius, P., Rasmussen, F. and Lundberg, I. (2008). Adolescent mental health predicts quitting smoking in adulthood: a longitudinal analysis. *European Journal of Public Health*, 18, 66-70.
- [42] Borzecki, A.M. and Lee, A. (2005). Do poor health behaviours affect health-related quality of life and healthcare utilization among veterans? *Journal Ambulatory Care Manage*, 28, 141-156.
- [43] Buckley, T.C., Mozley, S.L., Bedard, M.A., Dewulf, A.C. and Greif, J. (2004). Preventive health behaviours, physical morbidity, and health related role functioning impairment in veterans with post-traumatic stress disorder. *Military Medicine*, 169, 536-540.
- [44] Centres for Disease Control and Prevention. Youth Risk Behaviour Surveillance – United States. Surveillance Summaries, 2007. *MMWR* 2008;57(No. SS-4) (<http://www.cdc.gov/mmwr>).
- [45] The Information Centre, Lifestyles Statistics (2008). The health survey for England 2006. (<http://www.ic.nhs.uk>).

- [46] Pesa, J.A. (1998). The association between smoking and unhealthy behaviours among national sample of Mexican-American adolescents. *The Journal of School Health*, 68, 376-380.
- [47] National Centre for Social Research (2007). Smoking, drinking and drug use among young people in England in 2006 (<http://www.ic.nhs.uk>).
- [48] Williams, J.O., Bell, N.S. and Amoroso, P.J. (2002). Drinking and other risk taking behaviours of enlisted male soldiers in the US Army. *Work*, 18, 141-150.
- [49] Naval Health Research Centre (1998). Relationships Among Changes in Health Behaviours in a Six-Year U.S. Navy Cohort.
- [50] Beckham, J.C., Becker, M.E., Hamlett-Berry, K.W., Drury, P.D., Kang, H.K., Wiley, M.T., Calhoun, P.S., Moore, S.D., Bright, M.A. and McFail, M.E. (2008). Preliminary Findings from a Clinical Demonstration Project for Veterans Returning from Iraq or Afghanistan. *Military Medicine*, 173, 448-453.
- [51] Morgan, B.J. (2001). Evaluation of an educational intervention for military tobacco users. *Military Medicine*, 166, 1094-2004.
- [52] IOM (Institute of Medicine). 2009. *Combating Tobacco Use in Military and Veteran Populations*. New York, NY: The Guilford Press.
- [53] Peterson. L.P. (2006). Clinical health psychology and behavioural medicine in military healthcare settings. In C.H. Kennedy and E.A. Zillmer (Eds.), *Military psychology: Clinical and Operational Applications* (pp. 74-104). Washington DC: American Psychological Association.
- [54] Bray, R.M. and Brown, J.M. (2008). Influence of military leaders, units, and deployment on tobacco use. Paper presented at 50th Annual conference of the International Military Testing Association, Amsterdam 2008 (www.imta.org).
- [55] Bray, R.M., Pemberton, M.R., Williams, J., Green, K.J. and Hunter, C.M. (2008). Peer and role model influences on cigarette smoking. Paper presented at 50th Annual conference of the International Military Testing Association, Amsterdam 2008 (www.imta.org).
- [56] Joseph, A.M., Muggli, M., Pearson, K.C. and Lando, H. (2005). The Cigarette Manufacturers' Efforts to Promote Tobacco to the U.S. Military. *Military Medicine*, 170, 874-880.
- [57] Arvey, S.R. and Malone, R.E. (2008). Advance and Retreat: Tobacco control policy in the U.S. military. *Military Medicine*, 173, 985-991.
- [58] Unpublished survey.
- [59] Bech, P., Olsen, L.R., Kjoller, M. and Rasmussen, N.K. (2003). Measuring well-being rather than the absence of distress symptoms: a comparison of the SF-36 Mental Health subscale and the WHO-Five Well-Being Scale. *International Journal of Methods in Psychiatric Research*, 12, 85-91.
- [60] Hoffman, K.M., Haddock, C.K., Poston, W.S.C., Taylor, J.E., Lando, H.A. and Shelton, S. (2008). A formative examination of messages that discourage tobacco use among junior enlisted members of the United States military. *Nicotine & Tobacco Research*, 10, 653-661.