Chapter 6 – REFERENCES

[1] Barraquer, J.I. Keratomileusis. Int Surg. 1967;48:103-117.

[2] McDonald, M.B., Beuerman, R., Falzoni, W., Rivera, L. and Kaufman, H.E. Refractive surgery with the excimer laser. Am J Ophthalmol. 1987;103:469.

[3] Pallikaris, I.G., Papatzanaki, M.E., Stathi, E.Z., Frenschock, O. and Georgiadis, A. Laser in situ keratomileusis. Lasers Surg Med. 1990;10:463-468.

[4] Schallhorn, S.C., Blanton, C.L. and Kaupp, S.E. et al. Preliminary results of photorefractive keratectomy in active-duty United States Navy personnel. Ophthalmology. 1996;103:5-22.

[5] Clare, G., Pitts, J.A., Edgington, K. and Allan, B.D. From beach lifeguard to astronaut: occupational vision standards and the implications of refractive surgery. Br J Ophthalmol. 2010;94:400-405.

[6] Ambrosio, R. and Wilson, S. LASIK vs. LASEK vs PRK: advantages and indications. Semin Ophthalmol. 2003;18:2-10.

[7] Clare, G., Moore, T.C., Grills, C., Leccisotti, A., Moore, J.E. and Schallhorn, S. Early flap displacement after LASIK. Ophthalmology. 2011;118:1760-1765.

[8] Berjano, E.J., Navarro, E., Ribera, V., Gorris, J. and Alio, J.L. Radiofrequency heating of the cornea: an engineering review of electrodes and applicators. Open Biomed Eng J. 2007;1:71-76.

[9] Schwartz, A.R., Tinio, B.O., Esmail, F., Babayan, A., Naikoo, H.N. and Asbell, P.A. Ten-year follow-up of 360 degrees intrastromal corneal rings for myopia. J Refract Surg. 2006;22:878-883.

[10] Jung, S.W., Kim, M.J., Park, S.H. and Joo, C.K. Multifocal corneal ablation for hyperopic presbyopes. J Refract Surg. 2008;24:903-910.

[11] Fong, C.S. Refractive surgery: the future of perfect vision? Singapore Med J. 2007;48:709-18; quiz 719.

[12] Barsam, A. and Allan, B.D. Excimer laser refractive surgery versus phakic intraocular lenses for the correction of moderate to high myopia. Cochrane Database Syst Rev. 2012;1:CD007679.

[13] Alio, J.L., Ben-nun, J., Rodriguez-Prats, J.L. and Plaza, A.B. Visual and accommodative outcomes 1 year after implantation of an accommodating intraocular lens based on a new concept. J Cataract Refract Surg. 2009;35:1671-1678.

[14] Alio, J.L., Plaza-Puche, A.B., Javaloy, J., Ayala, M.J., Moreno, L.J. and Pinero, D.P. Comparison of a New Refractive Multifocal Intraocular Lens with an Inferior Segmental Near Add and a Diffractive Multifocal Intraocular Lens. Ophthalmology. 2012.

[15] Dennis, R.J., Beer, J.M., Baldwin, J.B., Ivan, D.J., Lorusso, F.J. and Thompson, W.T. Using the Freiburg Acuity and Contrast Test to measure visual performance in USAF personnel after PRK. Optom Vis Sci. 2004;81:516-524.

[16] Schallhorn, S.C., Tanzer, D.J., Kaupp, S.E., Brown, M. and Malady, S.E. Comparison of night driving performance after wavefront-guided and conventional LASIK for moderate myopia. Ophthalmology, 2009;106:702-709.

[17] Schallhorn, S.C., Farjo, A.A., Huang, D., Boxer Wachler, B.S., Trattler, W.B., Tanzer, D.J., Majmudar, P.A. and Sugar, A. Wavefront-guided LASIK for the correction of primary myopia and astigmatism. Ophthalmology, 2008;115:1249-1261.